

Microlensing, flattening, Ω_B and h : from assumptions, via data to tentative conclusions

Vesna Milošević-Zdjelar

*Dept. of Physics & Astronomy, University of Manitoba, Winnipeg, Mb,
R3T 2N2, Canada*

Srdjan Samurović and

*Dipartimento di Astronomia, Università degli Studi di Trieste, Via
Tiepolo 11, I-34131, Trieste, Italy*

Milan M. Ćirković¹

*Dept. of Physics & Astronomy, SUNY at Stony Brook, Stony Brook,
NY 11794-3800, USA*

Abstract. Microlensing observations can be used for determining the shape of the Milky Way's halo. It can be shown that the data are best described with moderately flattened halo, $0.5 \lesssim q \lesssim 0.6$. We discuss, by taking into account this result, the constraints on the baryonic mass-density parameter, Ω_B , and their implications on the value of the Hubble constant H_0 , i.e. h . Our conclusion is that current data, in order to satisfy BBNS and Galaxy dynamics constraints, strongly suggest that h should take values within the lower part of the permissible range, $h \sim 0.5$.

1. Microlensing and the flattening of the Galactic halo

We discuss constraints on the universal baryonic mass-density parameter $\Omega_B \equiv \rho_B/\rho_{\text{crit}} = 8\pi G\rho_B/3H_0^2$ and a Hubble parameter h ($H_0 = 100 h$ km/s/Mpc), using all available microlensing (ML) data and considering flattened Galactic dark halo. Since Ω_B and h are uncertain and correlated, we tried to break this degeneracy with the other cosmological information: measured optical depths from ML. In recent paper we examined the relation between galactic halo flattening and baryonic dark matter (Samurović, Ćirković & Milošević-Zdjelar 1999, SĆMZ), using mass density for flattened instead of spherical halo, and optical depth τ as a function of Galactic coordinates, derived by Sackett & Gould 1993 (eqs. 2.1 and 2.5). Comparing solutions for LMC, SMC, M31 and a source at 50 kpc with all available ML data, we obtained $0.5 \lesssim q \lesssim 0.6$ for halo flattening. Assuming that MACHOs have finite mass-to-light ratio (white or brown dwarfs), in SĆMZ we determined total mass of the MACHO halo as a function of flattening (eq. 9), and cosmological density parameter Ω_{MACHO} (eq. 10).

¹Affiliated to: Astronomska Opservatorija, Volgina 7, 11000 Belgrade, Serbia




Fig. 1: Dependence of Ω_B on h for flattening $q = 0.6$, truncation radius 50 kpc, MACHO halo mass within 50 kpc for $q = 0.6$. The sum of main baryonic components is Ω_{total} . High value $0.09 \lesssim \Omega_B \lesssim 0.1$, is within constraints of CMB observations: $\Omega_B < 0.28$ (Lineweaver et al. 1997), and recent BBNS and light element abundances constraints (Burles et al. 1999). At higher redshifts, high $\Omega_B > 0.017h^{-2}$ can account for observational results of Ly α forest. Our results fit into theoretical requirements of high resolution spectroscopy of the IGM (e.g. Burles & Tytler 1998).

2. Tentative conclusions

Recent estimate of total halo mass within 50 kpc is $\sim 5.4^{+0.2}_{-3.6} \times 10^{11} M_{\odot}$ (Wilkinson & Evans 1999). That can be obtained when we assume flattening parameter $q \sim 0.6$. Spherical halo has serious problems concerning BBNS limits.

Value $h \sim 0.5$ is in accord with $h = 0.39^{+0.14}_{-0.13}$ obtained by combining the baryon budget with CMBR measurements (Hannestad 1999). Realistic flattening and MACHO-dominance in the inner halo are incompatible with $h > 0.75$.

Acknowledgments. Authors acknowledge financial support of the Univ. of Trieste, and kind encouragement of Dr. Penny Sackett and Dr. Geza Gyuk.

References

- Burles, S., Nollett, K.M., Truran, J.N., & Turner, M.S. 1999, Phys. Rev. Lett., 82 (21), 4176
- Burles, S., & Tytler, D. 1998, ApJ, 499, 699
- Hannestad, S. 1999, MNRAS, in press, preprint astro-ph/9906246
- Lineweaver, C. H., Barbosa, D., Blanchard, A., & Bartlett, J. G. 1997, A& A, 322, 365
- Sackett, P. D., & Gould, A., 1993, ApJ, 419, 648
- Samurović, S., Ćirković, M. M., & Milošević-Zdjelar, V. 1999, MNRAS, in press (SĆMZ), preprint astro-ph/9907165
- Wilkinson, M.I., & Evans, N.W. 1999, MNRAS, in press, preprint astro-ph/9906197